

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

1. Induct on $|G|$.

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

1. Induct on $|G|$.
2. Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

1. Induct on $|G|$.
2. Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
3. Case 1: $p \nmid |C_i|$ for some i .

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

1. Induct on $|G|$.
2. Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
3. Case 1: $p \nmid |C_i|$ for some i .
4. Case 2: $p \mid |C_i|$ for all i .

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

1. Induct on $|G|$.
2. Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
3. Case 1: $p \nmid |C_i|$ for some i .
4. Case 2: $p \mid |C_i|$ for all i .
5. Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

1. Induct on $|G|$.
2. Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
3. Case 1: $p \nmid |C_i|$ for some i .
4. Case 2: $p \mid |C_i|$ for all i .
5. Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
6. The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

1. Induct on $|G|$.
2. Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
3. Case 1: $p \nmid |C_i|$ for some i .
4. Case 2: $p \mid |C_i|$ for all i .
5. Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
6. The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
7. The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

- ✓ Induct on $|G|$.
2. Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
3. Case 1: $p \nmid |C_i|$ for some i .
4. Case 2: $p \mid |C_i|$ for all i .
5. Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
6. The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
7. The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

- ✓ Induct on $|G|$.
- ✓ Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
- 3. Case 1: $p \nmid |C_i|$ for some i .
- 4. Case 2: $p \mid |C_i|$ for all i .
- 5. Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
- 6. The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
- 7. The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

- ✓ Induct on $|G|$.
- ✓ Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
- ✓ Case 1: $p \nmid |C_i|$ for some i .
- 4. Case 2: $p \mid |C_i|$ for all i .
- 5. Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
- 6. The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
- 7. The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

- ✓ Induct on $|G|$.
 - ✓ Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
 - ✓ Case 1: $p \nmid |C_i|$ for some i .
 - ✓ Case 2: $p \mid |C_i|$ for all i .
5. Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
 6. The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
 7. The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

- ✓ Induct on $|G|$.
- ✓ Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
- ✓ Case 1: $p \nmid |C_i|$ for some i .
- ✓ Case 2: $p \mid |C_i|$ for all i .
- ✓ Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
- 6. The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
- 7. The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

- ✓ Induct on $|G|$.
- ✓ Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
- ✓ Case 1: $p \nmid |C_i|$ for some i .
- ✓ Case 2: $p \mid |C_i|$ for all i .
- ✓ Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
- ✓ The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
- 7. The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).

Strategy (First Sylow Theorem)

Theorem

Let G be a finite group and p a prime such that $p^k \mid |G|$. Then G has a subgroup of order p^k .

- ✓ Induct on $|G|$.
- ✓ Apply the class equation in combination with $|C| = [G : C(a)]$, where $a \in C$, to count elements in G .
- ✓ Case 1: $p \nmid |C_i|$ for some i .
- ✓ Case 2: $p \mid |C_i|$ for all i .
- ✓ Now $p \mid |Z(G)|$, so $Z(G)$ has a (normal!) subgroup N of order p .
- ✓ The induction hypothesis applied to G/N gives $T \leq G/N$ with $|T| = p^{k-1}$.
- ✓ The pullback H of T (so $T = H/N$) is the subgroup we want (see Theorem 8.24).