

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

1. The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

1. The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).
2. The  $P$ -conjugates of the  $K_i$  are other  $K_i$  and partition  $S = \{K_1, \dots, K_t\}$  into sets of size  $p^k$  (using Theorem 9.25).

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

1. The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).
2. The  $P$ -conjugates of the  $K_i$  are other  $K_i$  and partition  $S = \{K_1, \dots, K_t\}$  into sets of size  $p^k$  (using Theorem 9.25).
3. One of the sets  $K_i$  must have size  $t_0$ .

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

1. The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).
2. The  $P$ -conjugates of the  $K_i$  are other  $K_i$  and partition  $S = \{K_1, \dots, K_t\}$  into sets of size  $p^k$  (using Theorem 9.25).
3. One of the sets  $K_i$  must have size  $t_0$ .
4. Lemma 9.26 implies  $P = K_i$ .

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

- ✓ The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).
2. The  $P$ -conjugates of the  $K_i$  are other  $K_i$  and partition  $S = \{K_1, \dots, K_t\}$  into sets of size  $p^k$  (using Theorem 9.25).
3. One of the sets  $K_i$  must have size  $t_0$ .
4. Lemma 9.26 implies  $P = K_i$ .

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

- ✓ The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).
- ✓ The  $P$ -conjugates of the  $K_i$  are other  $K_i$  and partition  $S = \{K_1, \dots, K_t\}$  into sets of size  $p^k$  (using Theorem 9.25).
- 3. One of the sets  $K_i$  must have size  $t_0$ .
- 4. Lemma 9.26 implies  $P = K_i$ .

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

- ✓ The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).
  - ✓ The  $P$ -conjugates of the  $K_i$  are other  $K_i$  and partition  $S = \{K_1, \dots, K_t\}$  into sets of size  $p^k$  (using Theorem 9.25).
  - ✓ One of the sets  $K_i$  must have size  $t_0$ .
4. Lemma 9.26 implies  $P = K_i$ .

# Strategy (Second Sylow Theorem)

## Theorem

*If  $P$  and  $K$  are Sylow  $p$ -subgroups of a group  $G$ , then there exists  $x \in G$  such that  $P = x^{-1}Kx$ .*

- ✓ The conjugates of  $K$  are  $K = K_1, \dots, K_t$ , with  $(p, t) = 1$  (using Theorem 9.25).
- ✓ The  $P$ -conjugates of the  $K_i$  are other  $K_i$  and partition  $S = \{K_1, \dots, K_t\}$  into sets of size  $p^k$  (using Theorem 9.25).
- ✓ One of the sets  $K_i$  must have size  $t_0$ .
- ✓ Lemma 9.26 implies  $P = K_i$ .