

MATH 142 Midterm Exam #1

October 6, 2006

NAME: _____

- No calculators are allowed on this exam.
- Answers such as $\frac{23.5}{30} - \frac{2^5}{3.34}$ are perfectly fine!! However you MUST simplify expressions such as $\sin(\pi/3)$.
- Please show all your work. You may use back pages if necessary. You may not receive full credit for a correct answer if there is no work shown.
- Please include all information about u-substitutions, and use correct mathematical grammar in the presentation of your solution.

Problem	Points	Score
1	20	
2	30	
3	15	
4	15	
5	20	
total	100	

$$\sum_{i=1}^n a = a \cdot n$$

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

1. Definition of the Integral. Recall the definition of the definite integral for a continuous function $f(x)$ on the interval $[a, b]$.

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$

Calculate the integral below using the definition of the integral.

$$\int_0^4 3 - x dx$$

- (a) First, find the following quantities:

$$\Delta x = \underline{\hspace{2cm}} \qquad x_i = \underline{\hspace{2cm}} \qquad f(x_i) = \underline{\hspace{2cm}}$$

- (b) Next, using the quantities above and the summation formulas on the front page of the exam, simplify $\sum_{i=1}^n f(x_i) \Delta x$ into an expression without the summation notation.

- (c) Last, evaluate the limit, $\lim_{n \rightarrow \infty} (\sum_{i=1}^n f(x_i) \Delta x)$.

Note: you can *check* your answer by using the Fundamental Theorem of Calculus.

2. Integrals. Evaluate the following definite and indefinite integrals.

(a) $\int (t^2 2^t - 7\sqrt{t} + 45t)t^{-2} dt$

(b) $\int_{\sqrt{\frac{\pi}{12}}}^{\sqrt{\frac{\pi}{2}}} x \sin(3x^2) dx$

(c) $\int_0^{4/3} \frac{3}{9x^2 + 16} dx$

3. Let $f(t)$ be the continuous function graphed below and let $g(x) = \int_{-3}^x f(t) dt$

(a) Evaluate $g(4)$.

(b) Find $g'(x)$ and evaluate $g'(0)$.

(c) Find the intervals over which $g(x)$ is concave up.

4. Evaluate the following limits. Show work and indicate if/when L'Hôpital's rule is used.

(a) $\lim_{x \rightarrow \infty} \frac{\tan^{-1}(x)}{\frac{6}{x} - \pi}$

(b) $\lim_{x \rightarrow 0^+} x^2 \ln(x^2 + x)$

5. Volume. Note: On this problem, you can earn partial credit for parts (b)-(e) by sketching an arbitrary slice of the volume in the space provided in the left margin.

Consider the region, R which is bounded between the curves

$$y = (x - 2)^2 \quad y = x + 4$$

- (a) Sketch the region and label the points of intersection.
- (b) Write an integral for the volume of the solid formed by rotating the region R about the x-axis. DO NOT evaluate the integral.
- (c) Write an integral for the volume of the solid formed by rotating the region R about the line $x = -2$. DO NOT evaluate the integral.
- (d) Now consider the solid whose base is the region R and whose cross-sections above the xy-plane and perpendicular to the x-axis (i.e. slices parallel to the y-axis) are squares. Write an integral for the volume of this solid, but DO NOT evaluate the integral.